



PCT/CH 2004/000504

REC'D 24 AUG 2004

WIPO PCT



INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

GB04/504

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

P. Mahoney

Dated 3 August 2004

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)


Patents Form 1/77

Patents Act 1977
(Rule 16)

1/77

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

1. Your reference

3008908

20AUG03 E831471-1 D10149

P01/7700 0.00-0319463.6

2. Patent application number
(The Patent Office will fill in this part)

0319463.6

20 AUG 2003

3. Full name, address and postcode of the or of each applicant (underline all surnames)Givaudan SA
Chemin de la Parfumerie 5
1214 Vernier
SwitzerlandPatents ADP Number (*if you know it*)

8408031001

If the applicant is a corporate body, give the country/state of its incorporation

Switzerland

4. Title of the Invention

COMPOUNDS

5. Name of your agent (*if you have one*)"Address for service" in the United Kingdom to which all correspondence should be sent (*including the postcode*)Centre for Innovative Technology (Givaudan UK Ltd.)
76-80 Church Street, Staines
Middlesex TW18 4XR
United KingdomPatents ADP number (*if you know it*)

8447815001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (*if you know it*) the or each application number

Country

Priority application number
(*if you know it*)Date of filing
(*day/month/year*)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(*day/month/year*)8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (*Answer 'Yes' if:*)

yes

a) any applicant named in part 3 is not an inventor, or
b) there is an inventor who is not named as an applicant, orc) any named applicant is a corporate body.
See note (d)

Patents Form 1/77

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 10

Claim(s) 1

Abstract 1

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77) 1

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date 20/08/03

12. Name and daytime telephone number of person to contact in the United Kingdom

Colin Brown (office time) Tel. No: 01/784417721

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505
 b) Write your answers in capital letters using black ink or you may type them.
 c) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
 d) If you have answered 'Yes' Patents Form 7/77 will need to be filed
 e) Once you have filled in the form you must remember to sign and date it.
 f) For details of the fee and ways to pay please contact the Patent Office

Patents Form 1/77

30089GB

DUPLICATE

1

COMPOUNDS

This invention related to a method of providing meat flavouring to foodstuffs and to novel compounds for providing such flavouring.


5

It is known that the formation of meat flavour proceeds via a sequence of steps known as the Maillard reaction. The sequence may be represented as follows:

The Maillard reaction is also known as non-enzymatic browning. The Amadori compounds are
10 key intermediates in the formation of flavours.

One amino acid for which no Amadori compound has hitherto been identified is cysteine ($\text{CH}_2(\text{SH})\text{CH}(\text{NH}_2)\text{COOH}$). Cysteine plays an important role in the development of flavour in meat, and this is believed to result from its ability to release hydrogen sulphide under the
15 conditions of the Maillard reaction (see, for example, Kobayashi and Fujimaki in Agric. Biol. Chem., 29, 698-81965). Numerous references have attributed to sulphur compounds a central role in the formation of meat flavours. However, unlike all other amino acids, cysteine has hitherto been believed to inhibit the Maillard reaction, because it tends to form relatively stable 2-glycosylthiazolidine-4-carboxylic acids, and this reaction competes with, and may even
20 prevent, the formation of Amadori compounds.

It has now been found, surprisingly, that, not only is it possible to make Amadori compounds from cysteine, but also that these compounds are particularly effective at conferring meat flavour in foodstuffs. The invention therefore provides a compound of the formula I:



25


I

in which R is hydrogen when $n=1-4$, or methyl when $n=0-3$.

30089GB

2

Although the compound is depicted as being of formula I, this formula is in fact in equilibrium with two other forms, Ia and Ib, as shown below:



- 5 These cyclic forms are included within the scope of the invention. Also included are all other isomeric forms, of which a number are possible (for example, when L-cysteine reacts with a D- or L-sugar).

The compounds may be prepared by known methods. Amino acids or their salts may be reacted
 10 with an aldose either in the presence of about 20% of water (see E.F.L.J. Anet, Austr. J. Chem. 10, 193-197 (1957)) or in an organic solvent (see K. Heyns and H. Paulen, Liebigs Ann. Chem., 622, 160 (1959)). In all preparations of Amadori compounds so far known, a reaction medium with low water content is essential to bring about the required condensation with aldoses to glycosyl amines. However, the preparation of the Amadori compounds of cysteine can also
 15 conveniently be carried out in dilute aqueous solutions or in mixtures of water with water-miscible solvents. It is not necessary to isolate the initially-formed 2-glycosylthiazolidine-4-carboxylic acids. The formation of the Amadori compounds from cysteine plus sugar or from 2-glycosylthiazolidine-4-carboxylic acid is subject to general acid catalysis. For this reason, the reactions are preferably carried out in the presence of buffers having a pH of between 1 and 9,
 20 preferably between 2 and 6.

30089GB

3

Although the Amadori rearrangement proceeds at room temperature, it is preferred to accelerate it by heating. The preferred temperature will depend on the solvents used and on the reactant concentrations, but the temperature generally lies between 60° and 110°C. At 90°-100°C, the preferred reaction time of the aqueous buffered reaction mixture is from 1-12 hours.

5

The sugars to be used for the present invention are aldoses with 3-6 carbon atoms, including aldopentoses, aldohexoses and deoxyaldohexoses. Specific sugar examples are the aldopentoses ribose, arabinose, xylose and lyxose, the aldohexoses glucose, mannose and galactose and the 6-deoxyaldohexoses rhamnose, fucose and chinovose. Mixtures thereof can also be used. In addition, all stereoisomeric forms of these sugars may be used.

10

The cysteine may be used as the compound itself or in the form of a derivative or a salt, which form free cysteine under the reaction conditions. Both L- and D-cysteine and mixtures thereof may be used. The molar proportions of the cysteine and the aldose may vary between 2:1 and 1:5, preferably between 1.2:1 and 1:1.5.

15

The buffers used may comprise mixtures of weak acids and bases or compounds that possess both an acidic and a basic function. Suitable acids include carboxylic acids and phosphoric acid. Typical examples of bases include the salts of the previously-mentioned acids, for example, their alkali metal salts of their salts with nitrogen-containing bases such as pyridine, α -picoline and tertiary amines. Primary and secondary amines may also be used, but preferably only if the reaction medium contains a large excess of water, to prevent the competitive reactions of these amines with the aldose to glycosylamines and Amadori rearrangement compounds of these amines. Examples of compounds with both acidic and basic functions are amino acids or their salts, for example glycine, proline and monosodium glutamate, or ribonucleotides or their salts, such as disodium inosine-5'-monophosphate. The buffers are preferably used in concentrations of from 0.1 to 10 times the molar concentration of cysteine.

20

25

The Amadori compounds thus prepared may be isolated as crude products and purified by known methods to give the compounds in colourless, crystalline form.

30

30089GB

4

The Amadori compounds of this invention may be used to impart a meat-like taste to foodstuffs, both meat and non-meat, by incorporation of the compounds or compositions containing them into the foodstuff and subsequent heating. The preferred heating time depends on the water activity, the temperature used and the nature of the Amadori compound, but typically it is about 5 the time needed to fry or cook meat.

The concentration required may vary over a wide range, depending on the desired taste and aroma effects, but as a general guide, they are employed in concentrations of from 5 to 5000 ppm by weight of foodstuff in consumable form. Good results are obtained by using from 200-10 2000 ppm.

The Amadori compounds of this invention may be used in combination with other known food additives, such as flavour compounds, aroma enhancers, and the like.

15 The invention is further described with reference to the following non-limiting examples.

EXAMPLE I

Preparation of Amadori compound of L-cysteine and D-xylose

20 A solution of 60.5 g (0.50 mol) L-cysteine 82.5 g (0.55 mol) D-xylose and 50 g (about 0.60 mol) of a mixture of methylated pyridines in 750 ml demineralised water was heated to about 95°C on a water bath in a 1L round-bottomed flask equipped with a reflux condenser. After 0.5 h, 120 g (2.0 mol) acetic acid was added and the solution was heated for a further 6 h at 90°C. The conversion to Amadori compound was then about 25%. The solution was cooled to room 25 temperature and the solution was then continuously extracted with ether to remove the methylated pyridines. The solution was passed through a 40 x 4 cm column of Bio-Rad AG 50W-X4 (H⁺) cation exchange resin. The column was washed with 1 L water and the absorbed substances were displaced from the column with 0.3N ammonia. Fractions containing more than 50% of the compound were combined and evaporated *in vacuo* to small volume. After the

30089GB

5

addition of just enough ethanol to cause no cloudiness, crystalline compound separated. The pure compound did not melt but decomposed above 170°C.

Found : C 37.87; H 5.96; N 5.51; S 12.65; O 37.80.

Calc. for $C_8H_{15}NSO_6$ (253.27) : C 37.94; H 5.97; N 5.53; S 12.66; O 37.90.

5 The product was dried prior to element analysis (*in vacuo* over P_2O_5 , 18 h at 69°C).

The results of a ^{13}C NMR spectrum (in D_2O ; δ in ppm, referenced to sodium 3-trimethylsilyl [2,2,3,3- 2H_4]-propionate) were as follows:

10	<u>Major isomer</u>	<u>Minor isomer</u>	<u>assignment</u>
	26.9(t)	25.6(t)	C6
	53.8(t)	50.2(t)	C3
	61.5(d)	57.0(d)	C5
	65.2(t)	65.2(t)	CH ₂ OH
15	71.9(d)	71.6(d)	} CHOH
	75.0(d)	76.6(d)	
	81.5(s)	80.9(s)	C2
	173.4(s)	173.0(s)	carboxyl carbon

20 EXAMPLE 2

Preparation of Amadori compound of L-cysteine and D-xylose

L-cysteine.HCl monohydrate (17.6 g; 0.1 mol) was dissolved in 50 ml of demineralized water contained in a 150-ml round-bottomed flask. Disodium hydrogen phosphate (14.2 g; 0.10 mol) or 90% lactic acid (10 g; 0.10 mol) was added and the pH of the resulting solution adjusted to 5.0 by slowly adding with stirring an aqueous 50% sodium hydroxide solution. The sugar (0.11 mol) was then added and the mixture stirred for 1 h at 50 °C to obtain an equilibrium mixture of thiazolidine, cysteine and sugar. After cooling to room temperature, the pH was re-adjusted to 5.4 by addition of 50% sodium hydroxide solution. The water content was adjusted by

30 evaporation *in vacuo* to 30%. The flask was then equipped, via a double neck adapter, with a

30089GB

6

mechanical stirrer and a reflux condenser, and heated with stirring for 7 h in a thermostated water bath of 80 °C. The conversion to Amadori compound is 35%. Isolation was performed as described in Example 1.

5 EXAMPLE 3

Continuous process for production of Amadori compound of L-cysteine and D-xylose

The continuous reactor consisted of a 45x2.5 ID cm glass column with temperature control jacket packed with Bio-Rex 9 anion exchange resin (OH⁻), 100-200 mesh. A restriction
10 consisting of a 500 x 0.3 mm ID TeflonTM tubing was connected to the column outlet to prevent the formation of gas bubbles in the column. The column was thermostated at 90 °C. A 50% thiazolidine solution, prepared by heating equimolar amounts of cysteine and xylose in water for 1 h at 50 °C, was pumped through the column at a rate of 0.5 ml/min. Analysis of samples taken
15 at regular time intervals showed that the conversion to Amadori compound increased from 12% after 2 hrs to 35% after 6 hrs. The yield of the corresponding batch process was 22.5%. The Amadori compound was isolated from the column eluate by *evaporation in vacuo* and crystallization of the residue from water-ethanol as in Example 1.

EXAMPLE 4

20 Preparation of the Amadori compound of L-cysteine and D-ribose

The same equipment was used as was used in Example 1. 12.1g (0.1 mol) of L-cysteine, 15g (0.1 mol) of D-ribose and 27.2g (0.2 mol) of potassium dihydrogen phosphate in 150 ml demineralised water were heated for 6h at ca. 90°C under nitrogen. The resulting brown solution
25 was passed through a 35 x 2.5cm column of DowexTM 50W-X4 (H⁺) cation exchange resin. The neutral components were washed from the column with 1L water and the nitrogen-containing substances were displaced from the column with 0.3N ammonia. Fractions containing more than 50% of the compound were combined and evaporated *in vacuo* to small volume. After the addition of just enough ethanol to cause no cloudiness, crystalline compound separated. The

30089GB

7

pure compound did not have a sharp melting point but decomposed at 132°-135°C. The results of a ^{13}C NMR spectrum (in D_2O ; δ in ppm) were as follows:

	<u>Major isomer</u>	<u>Minor isomer</u>	<u>assignment</u>
5	26.8(t)	25.7(t)	C6
	55.2(t)	51.1(t)	C3
	61.9(d)	57.3(d)	C5
	65.6(t)	65.3(t)	CH_2OH
	74.0(d)	73.4(d)	} CHOH
10	75.9(d)	77.8(d)	
	82.7(s)	81.1(s)	C2
	174.0(s)	173.5(s)	carboxyl carbon

EXAMPLE 5

15 Preparation of the Amadori compound of L-cysteine and L-arabinose

The process of Example 3 was repeated with L-arabinose instead of D-xylose. The resulting crystalline material melted with decomposition at 171-173 °C.

The results of a ^{13}C NMR spectrum (in D_2O ; δ in ppm) were as follows:

20

	<u>Major isomer</u>	<u>Minor isomer</u>	<u>assignment</u>
	26.9(t)	25.8(t)	C6
	54.9(t)	51.7(t)	C3
	62.1(d)	57.2(d)	C5
25	65.4(t)	65.6(t)	CH_2OH
	73.7(d)	73.9(d)	} CHOH
	77.7(d)	76.2(d)	
	81.1(s)	82.8(s)	C2
	174.0(s)	173.6(s)	carboxyl group

30

30089GB

8

EXAMPLE 6**Preparation of the Amadori compound of L-cysteine and D-glucose**

An aqueous solution of 12.1g (0.1 mol) of L-cysteine, 19.8g (0.1 mol) of D-glucose and 11.5g (0.1 mol) of L-proline was concentrated *in vacuo* to a syrup containing approximately 20% of water. The syrupy mixture was heated on a water bath at 95°C for 2.5h. HPLC analysis showed that, in addition to the Amadori compound of L-cysteine, the Amadori compound of L-proline was also formed, in the ratio of 3:1. The brown reaction mixture was dissolved in 500ml water and the Amadori compounds recovered as described in Example 2. The results of a ¹³C NMR spectrum (in D₂O; δ in ppm) were as follows:

	<u>Major isomer</u>	<u>Minor isomer</u>	<u>assignment</u>
	27.1(t)	25.9(t)	C6
	54.1(t)	50.3(t)	C3
15	61.6(d)	57.2(d)	C5
	65.1(t)	65.1(t)	C6'
	63.3(t)	63.0(t)	CH ₂ OH
	71.6(d)	71.3(d)	} CHO
	73.3(d)	73.5(d)	
20	74.4(d)	76.1(d)	} CHO
	81.8(s)	81.1(s)	
	173.6(s)	173.0(s)	C2
			carboxyl carbon

EXAMPLE 7**25 Preparation of Amadori compound of L-cysteine and L-rhamnose**

A solution of 3.3 g (0.30 mol) L-cysteine 49g (0.27 mol) L-rhamnose monohydrate and 30 g (about 0.0 mol) of a mixture of methylated pyridines and 72g (1.20 mol) acetic acid in 450 ml demineralised water was heated for 8h on a steam bath in a 1L round-bottomed flask equipped with a reflux condenser. The resulting brown reaction mixture was purified as described in

30089GB

9

Example 1. The resulting crystalline compound decomposed at 183°C. The results of a ^{13}C NMR spectrum (in D_2O ; δ in ppm) were as follows:

	<u>Major isomer</u>	<u>Minor isomer</u>	<u>assignment</u>
5	21.1 or 21.2(q)	21.1 or 21.2(q)	CH_3
	26.9(t)	25.9(t)	C6
	54.3(t)	50.8(t)	C3
	62.1(d)	57.0(d)	C5
	69.9(d)	69.7(d)	
10	75.2(d)	74.7(d)	CHOH
	75.8(d)	75.4(d)	
	81.3(s)	81.8(s)	C2
	173.9(s)	173.5(s)	carboxyl carbon

15 EXAMPLE 8

Sensory evaluation of Amadori compounds

A beef broth was prepared. Four portions of this broth were treated as follows:

- 20 Portion A: no treatment (control);
 Portion B: addition of 480ppm (4.0mmol/L) L-cysteine + 552ppm (4.0 mmol/L) D-xylose;
 Portion C: addition of 1000ppm (4.0 mmol/L) 2-xylosylthiazolidine-4-carboxylic acid
 (prepared according to R. Bognár *et al*, Liebigs Ann.Chem.738, 68 (1970);
 Portion D: addition of 1000ppm (4.0 mmol/L) of the Amadori compound of Example 1.

25

The mixtures were bottled and sterilised in an autoclave for 1h at 100°C. On sensory evaluation, Mixture D was preferred, because of its much stronger meaty taste. Moreover, it lacked the cysteine-like off-taste of Mixture B and the hydrogen sulphide-like off-odour of Mixture C.

30

30089GB

10

EXAMPLE 9**Test in a gravy**

A gravy was prepared by heating 100g of commercial, flavoured margarine to 160°C. On
5 reaching this temperature, 35ml water was added. This was used as an unflavoured control.

A second gravy was prepared by adding 100mg (1000ppm) of the Amadori compound of
Example 2 to 100g of margarine and heating to 160°C. When this temperature was reached,
35ml water was added.

10


A taste comparison revealed a considerable preferment for the gravy containing the Amadori
compound, because of its pronounced, nicely roast, meaty taste.

30089GB

11

CLAIMS

1. A compound of the formula I:



5

in which R is hydrogen when n=1-4, or methyl when n=0-3.


2. A method of preparing a compound according to claim 1, comprising the reaction of cysteine with a sugar in either aqueous solution or in a mixture of water and water-miscible solvents.
3. A method of conferring a meaty flavour or aroma on a foodstuff, comprising the addition to the foodstuff before heating of a compound according to claim 1.
4. Use of a compound according to claim 1 to confer on a foodstuff a meaty flavour or aroma.

30089GB

12

ABSTRACT

The Amadori compounds of cysteine



5

in which R is hydrogen when n=1-4, or methyl when n=0-3. The compounds are useful for conferring meaty tastes and aromas on foodstuffs.